


# Canada's Fraser River

## Reasons for sockeye salmon declines with a comparison to Bristol Bay

Seminars and Workshops

Dr. Garry R. Wobdy, Ph.D.  
Fisheries Resource Management Consulting  
[www.fishresource.com](http://www.fishresource.com)



22

2011

01

## □Ø

### Table of □Ø Content§

Introduction .....	1 □Ø
Fraser River freshwater environment .....	3 □Ø
Contaminants .....	3□Ø
Land Use .....	4□Ø
Mining .....	4□Ø
Hydroelectric .....	4□Ø
Urbanization .....	5□Ø
Forestry .....	5□Ø
Agriculture .....	5□Ø
Predation .....	5□Ø
Climate Change .....	5□Ø
Marine environment near the Fraser River .....	6 □Ø
Contaminants .....	6□Ø
Land/Marine Waters Use .....	7□Ø
Predation .....	7□Ø
Climate Change .....	7□Ø
Fisheries Management .....	8 □Ø
Fraser River Management .....	8□Ø
Bristol Bay Management .....	9□Ø
Influences on the Bristol and Fraser River sockeye .....	10□Ø
Habitat .....	11□Ø
Aquaculture .....	11□Ø
Human development .....	12□Ø
Cumulative impacts .....	12□Ø
Conclusions .....	13□Ø
References .....	14□Ø

□δ

□δ

□δ

## Canada's Fraser River: □δ Reasons for sockeye salmon declines □δ□δ with a □δ comparison to Bay □δ δ

□δ

Sarah O'Neal □δ

Dr. Carol Ann Woody □δ

Fisheries Research and Consulting □δ

[www.fish4thfuture.com](http://www.fish4thfuture.com) □δ

□δ

### Introduction □δ

□δ

Recently, Fraser River sockeye populations have been compared to those of Bay by proponents of mining projects in Bay who cite Fraser River sockeye as an example of 'the difference' between mining and fisheries (Johng 2001). Due to distinct physical and biological nature, as well as different levels of urbanization and industrialization, the two systems make an unlikely comparison. However, current state of the Fraser River system with impaired water quality, human development, changes in fish habitat and prey bases, and climate change has resulted in low productivity of the Fraser river sockeye in fifty years. □δ □δ

□δ

Fraser River sockeye salmon populations are suffering from myriad problems associated with urban and industrial development, leading to dramatic increases in productivity, multiple fisheries closures, and federal and international population listings. In freshwater, there are stressors from contamination (from mining, wood products and other industrial facilities), introduced predators, and increased river temperatures. In estuarine and marine environment, stressors are related to house and industrial waste, shipping, loss of habitat, warmer marine water temperatures. □δ

□δ

While the blame for declines simply cannot be pointed single direction, the current state of Fraser sockeye is an unfortunate and disastrous example of the coexistence of human development and salmon. □δ


□δ

### Status of the Fraser salmon □δ

□δ

The Fraser River is known as the greatest salmon river in the world and indeed is Canada's largest salmon producer (Bürgner 1991). □δ Sockeye (Oncorhynchus nerka) □δ

are the most commercially valuable species in the Fraser, and generated hundreds of millions of dollars annually until the mid-1990s (DFO 2001). In recent decades, however, the total returns declined productivity (recruit per spawner), and commercial values suffered wide fluctuations, and ultimately significant declines (Pacific Salmon Commission data, 2011). Productivity currently stands at 1.25 million, indicating populations are barely replacing themselves (Figure 1, Peterman 2010). Returns resulted in the fisheries closures in 2007 and 2008, including three consecutive years from 2007 to 2009, when total runs failed to exceed 1 million fish (Figure 1, Pacific Salmon Commission data).



**Figure 1. Total Fraser River sockeye returns (top) = total harvest + escapement from year to year moving average of total returns per spawner across all Fraser River sockeye stocks (not including the minor jacks component) divided by total spawners 4 years before. The horizontal dashed line indicates the productivity at which population can replace itself, i.e. returns/spawner = 1.25. Pacific Salmon Commission data in Peterman 2010.**

During the most recent conservation status review, the International Union for the Conservation of Nature (IUCN) categorized 50% of Fraser River sockeye salmon stocks as threatened: one Critically Endangered, three as Endangered, and 16 Vulnerable

(Rand 2008). □**□** Chinook salmon monitoring other fish are designated endangered □**□** by □**□** Canadian government Committee on the Status of Endangered Wildlife in Canada □**□** (COSWEIC; DF 2001). □**□**

□**□** Prodigious research into causes of decline includes an ongoing \$20 million dollar □**□** federal judicial inquiry. □**□** Data results suggest salmon and their essential habitats suffer □**□** from a multitude of stressors. □**□** Following discussion summarizes some of the likely □**□** reviewed and gray literature on the Fraser River sockeye declines, and concludes with a □**□** brief comparison of the Fraser River with the world's largest salmon producing □**□** system, the Bristol Bay watershed in Alaska. □**□** A discussion will be □**□** updated □**□** results emerge from the Columbia Commission federal inquiry, currently underway. □**□**

## Fraser River freshwater environment □**□**

### □**□** Contaminants □**□**

□**□** MacDonald et al. (2011) systematically evaluated over 200 aquatic contaminants in the □**□** Fraser River basin in addition to potential anadromous salmonids. □**□** The □**□** study indicates contaminated surface water sediment, as well as accumulation of □**□** contaminants in fish, could pose hazards to the spawning and migrating salmon. □**□** Primary elements of concern include total suspended solids (TSS), turbidity, nutrients □**□** (nitrate, nitrite, and phosphorus), major ions (chloride, fluoride, and sulfate), metals □**□** (aluminum, arsenic, boron, cadmium, chromium, cobalt, copper, iron, lead, mercury, nickel, □**□** selenium, and silver), and phenols. □**□** Concentrations of 23,784 chlorodibenzene equivalents □**□** occurred at concentrations that may adversely affect □**□** reproduction. □**□** Data is insufficient to thoroughly examine impacts of □**□** endocrine □**□** disrupting chemicals such as pharmaceutical personal care products, industrial chemicals, □**□** pesticides, inorganic and organometallic compounds, and biogenic compounds (MPCA □**□** 2008), though authors concluded they undoubtedly were entering the Fraser River and □**□** likely have impacts on the sockeye salmon and reproduction. □**□** For example, the □**□** occurrence of female sterile salmon (MacDonald et al. 2011) is likely result of □**□** exposure to endocrine disruptors. □**□**

□**□** Sources of contamination numerous. Twenty-eight major mines (Figure 2), many small □**□** placer mines, 100 paper mills, 99 sawmills, wood mills, □**□** and other wood product □**□** facilities, 150 wastewater treatment facilities, 170 cement concrete facilities, 380 seafood □**□** processing facilities, 370 municipal wastewater treatment plants, 370 salterns □**□** facilities (Appendix 1), 830 municipal landfills, several manufacturing □**□** facilities, as well as a large industry operate within the watershed (MacDonald et al. □**□** 2011). Many of the aforementioned facilities are permitted to discharge pollutants of □**□** concern (MacDonald et al. 2011) and accidents occur. □**□** For instance, 510 spills reported □**□** from various facilities during a four-year period in 2010 (MacDonald et al. 2011). □**□** Of the □**□** 2866 sites listed in the Canadian Contaminated Sites Registry nearly 1500 years, 2600 (91%) □**□** were located in the Fraser River watershed (MacDonald et al. 2011). □**□** The number of □**□** contaminated sites is currently estimated to be approximately 5000 (MacDonald et al. 2011). □**□**

□

Human activities also contribute non-point source pollution to the Fraser River. □ Forest management activities, agricultural operations, and stormwater runoff can contain □ sediment, fertilizers, insecticides, fire retardants, and other contaminants (MacDonald et al. 2011, Nelitz et al. 2011). MacDonald (2011) indicate substantial quantities of □ suspended solids, □ nutrients, phenols, and total hydrocarbons have been released to the Fraser River from non-point sources. □ Finally, other sources of □ pollutants include persistent organic pollutants and mercury, which can also impact aquatic ecosystems (Mair et al. 2005), include forest fires, volcanoes, and carbon emissions (MacDonald et al. 2011).

□

## Land Use □

□

Nelitz et al. (2011) additionally examined impacts of □ mining, agriculture, □ hydroelectricity, urbanization, and water use on the freshwater environment and their potential impacts on □ Fraser sockeye salmon populations. □

□

### Mining □

Several types of □ mining take place in the Fraser Basin (Figure 2): □ planning, gravel mining, industrial mineral production, metal mining, oil and gas production, and coal mining. □ At one operating mine, Gibraltar, produces acid mine drainage associated with high levels of □ dissolved metals, exceeding federal and provincial effluent discharge criteria by □ several orders of magnitude (Ferguson 1987). □ Planning is □ the dominant mining activity in the basin and may have the most significant impacts to □ salmon due to □ sediment effects (Nelitz et al. 2011).




Figure 2. Distribution of □ different categories of □ large mines in the Fraser River Basin. □ Nelitz et al. 2011 □

□

□ effluent discharge criteria by □ several orders of □ magnitude (Ferguson 1987). □ Planning is □ the dominant mining activity in the basin and may have the most significant impacts to □ salmon due to □ sediment effects (Nelitz et al. 2011).

□

### Hydroelectric □

Two large hydroelectric projects within the basin, □ the George/Seton River power project and Alcan's Kemano Project on the Nechako River that affect water temperature and flow, at □ the time limiting migration ability of □ the sockeye salmon (Nelitz et al. 2011). □ The scale model hydropower projects, which can affect Total Gas Pressure (usually nitrogen supersaturation), gravel supply, and water temperature, also exist in the basin, though in □ the flywheel numbers (about 20) (Nelitz et al. 2011).

## **Urbanization**

Water demand was associated with high human densities, largely in lower portions of the Fraser River basin. Population growth, associated with urbanization, was 81% in the lower Fraser and 25% in municipalities south of the British Columbia in the past 20 years. Urbanization led to alterations to the saltation that from impervious surfaces including roads, changes in the hydrology, stream crossings and channelization.

## **Forestry**

While forest harvest has decreased significantly in recent decades there is more than one stream crossing per square kilometer in spawning areas and migration corridors (MOE 2008). Roads often serve as barriers to movement (Warren and Pandew 1998), and impact the fishery of anadromous salmonids (Groot and Margolis, 1996). Further, up to 90% of the streams in some watersheds were disturbed by Mountain Beetle infestation, potentially increasing fire risk and sedimentation as well as impacting the hydrology (Nelitzel et al 2011).

## **Agriculture**

The land area occupied by agriculture increased in the past twenty years. Agriculture can cause physical alteration to the streambank zones, and floodplains; increase sedimentation and destabilize stream banks causing widening of the streambed; remove vegetation which can increase stream temperatures; compact soil subsequently increasing runoff; deplete groundwater sources important to the maintenance of streams and temperature regimes; increase biochemical oxygen demand; introduce pathogens; and increase sedimentation, nutrients and contaminants through applications of manure, fertilizers, and pesticides.

## **Predation**

Predation of sockeye salmon occurs in freshwater and marine environments. Christensen and Triðes (2011) reviewed available literature on predation. Smallmouth bass as well as yellow perch are introduced species in the watershed, and are known to feed on salmon species, but little data exists regarding their impact on sockeye (Christensen and Triðes 2011). Hatchery wild salmon both compete with and directly prey upon sockeye (Appendix K, 2009, Tataran and Benjikian 2011, Ruggerone et al. 2011), although impacts are not well documented in the Fraser Basin. Salmon enhancement facilities in the Fraser River Basin are listed below in Appendix K. In addition to predation, hatchery fisheries are a source of potential contamination and have additional negative ecological effects on salmon populations (see below, Bubble et al. 2009).

## **Climate Change**

In British Columbia, minimum temperatures have increased 0.17°C per decade and precipitation has increased by 22% per century (Hinch and Martins 2011). Climate change has already caused

earlier snowmelt in British Columbia rivers (Stewart et al. 2005), and water temperatures in the Fraser River have increased at a rate of 0.33°C per decade, increasing overall water temperature by about 2°C in the past 60 years (Chittendon et al. 2009). Lakes in the region are also warming, altering timing of spring ice break-up and lake turnover (Schindler et al. 2005).

Temperature related factors have also received a great deal of attention with respect to a marked increase in mortality during river migration and on spawning grounds (Hinch and Martins 2011).

- x **Eggs.** Although sufficient data is lacking to thoroughly examine potential impacts of increased rainfall resulting from climate change, it is possible that increased rainfall is causing increased scour of redds, thereby decreasing overall egg survival (Hinch and Martins 2011).
- x **Fry.** Temperature increases may be facilitating increased predation on lake-rearing sockeye fry (Hinch and Martins 2011).
- x **Adult migrants.** Warmer river temperatures appear to decrease survival of adult migrants, particularly in early-run stocks, likely from a combination of exposure to temperatures above the 18°C thermal tolerance, increased energy required for migration at higher flows, and combined higher metabolism in elevated temperatures (Eliason et al. 2011, Hinch and Martins 2011). Pathogens including *Parvicapsula minibicornis* also develop more quickly in warmer temperatures (Cooke et al. 2004, Crossin et al. 2009), increasing physiological stress and decreasing swimming performance of adult migrants (Bradford et al. 2010, Wagner et al. 2005). Earlier migration timing, likely related to elevated temperatures, has coincided with en route and pre-spawning mortality exceeding 90% in some years, impacting larger stocks and pushing already threatened stocks such as Cultus Lake to near extinction (Cooke et al. 2004). These trends are expected to increase as climate change progresses (Hague et al. 2011, Rand et al. 2006).

## □ δ Marine environment near the Fraser River □ δ

### □ δ *Contaminants* □ δ

The Strait of Georgia borders British Columbia's main population centers of Vancouver and Victoria. □ δ About 90% of the marine pollution estimated through □ δ disposal of land-based waste from land-based activities (MOE 2006). □ δ Households □ δ generate about one third of what is from industrial sources (MOE □ δ 2006). □ δ Despite increases in the populations of those (Johannes et al. 2011) □ δ contaminants in the Strait of Georgia and general improvement in recent Best □ δ management practices including recycling programs and secondary or □ δ better treatment have improved in recent years (Johannes 2011) □ δ However, lead, □ δ polychlorinated biphenyls (PCBs), mercury, dioxins and furans, tri- □ δ butyltin were □ δ documented at higher concentrations in the waters, among birds and □ δ other biota in the Strait of Georgia (Johannes 2011). □ δ Pulp paper mills along the shores of the Strait were a major contributor of contaminants at least the 1980's when effluent treatment improved. □ δ In decades, polybrominated diphenylethers, personal care products and pharmaceuticals increased in the Strait of Georgia (Johannes et al. 2011). □ δ

□ δ

## **Land/Marine Waters Use □ δ**

□ δ

Shipping and marine vessels transport most goods and services across the coast, and may be source of noise, contaminants, accidental spills, and non-native species introductions through ballast water exchange, though Johannes et al. (2011) concluded that marine traffic has only limited direct interaction with sockeye habitat. Dredging lowered the main navigation channel at the mouth of the Fraser River kilometers over the past 30 years, though dredging activities are limited to periods when sockeye salmon are not in the estuary (EREMP 2006). Arteries have been removed throughout the lower Fraser River estuary, causing an estimated 40% habitat loss (Ellis et al. 2004) though although construction has slowed in recent decades, some have been removed to restore natural habitat (Johannes et al. 2011). □ δ

□ δ

## **Predation □ δ**

□ δ

Significant marine predators of Fraser sockeye salmon may include spiny dogfish (*Squalus acanthias*), sandsharks (*Cephaloscyllium laticeps*), and sand tiger sharks (*Anatopterus nictipinnis*; Christensen and Triðes 2001). Harbor seals (*Phoca vitulina richardsi*) and sea lions (*Zalophus californianus*) are also common predators and have increased dramatically since their protection in 1972 under the Fisheries Act (Forrest et al. 2009, Christensen and Triðes 2001). Pacific herring (*Clupea harengus pallasi*) and sandlances (*Ammodytes macrocephalus*) are unlikely to prey on sockeye salmon smolts, though they are likely competitors for food in the Strait of Georgia and have been increasing in numbers recent decades (Christensen and Triðes 2001). □ δ

□ δ

Many non-native introduced species in the Strait of Georgia prey upon and/or compete with Fraser River sockeye salmon. Though an estimated 10 introduced species, more than twice the number found throughout the remainder of the Canadian coast, a result of human population growth, aquaculture, and shipping activities (Johannes et al. 2011). Available data is inconclusive; recently documented Humboldt squid (*Dosidicus gigas*, Coërgrove 2005) may be significant predators of Fraser sockeye smolts (Christensen and Triðes 2001). □ δ

□ δ

Christensen and Triðes (2001) indicate that insufficient data exists to adequately identify key predators of sockeye salmon and the overall impact, as well understand the critical cumulative impact of predation on Fraser sockeye salmon and freshwater environments. □ δ

□ δ

## **Climate Change □ δ**

□ δ


Major cycles associated with climate and sea surface temperature in the North Pacific Ocean, the El Niño / South Oscillation (ENSO) and the Pacific Decadal Oscillation (PDO) have exhibited pattern changes in recent decades (Beamish 1999, Mantua et al. 1997). The Marine habitat for Fraser River sockeye salmon, the Strait of Georgia and the Gulf of Alaska, □ δ

Chittenden et al. 2019 found that the number of days with surface water temperatures above 18°C has increased from the late 1980's to the present, experienced warmer conditions than those during the previous period starting in the 1940s (Figure 3). The surface temperature (SST) has increased by about 1.5 °C over the last 60 years (Chittenden et al. 2019). Warmer temperatures are coincident with blooms of harmful algae *Heterosigma*. *Heterosigma* blooms can cause salmon mortality through diminished respiratory function and ability to uptake oxygen (Rensel et al. 2010).

**Figure 3.** Time series of monthly surface temperature anomalies from the long-term mean from 1986–2014 at Grouse Entrance Island, central Strait of Georgia (Johannes et al. 2011).

Temperature changes may also be due to causing a decline in zooplankton prey quality source for sockeye salmon (Figure 4; Johannes 2011). Declines in prey plankton coincide with increases of other species which sockeye may prey upon, although their food quality is considerably less fat content (Eloranta-Saharavi et al. 2019) and Ovalle, Johannes et al. (2011) conclude that warming temperatures coincide with declining food availability and quality, which may limit sockeye growth and decrease condition.

**Figure 4.** Declining zooplankton (*Neocalanus* sp.) abundance in the Strait of Georgia (Johannes and MacDonald 2009).



## Fisheries Management

### Fraser River Management

Management of Fraser sockeye and other salmon falls under myriad legal statutes. Due to the international range of the marine ecosystem, the fishery is subject to

international Pacific Salmon Treaty between the U.S. and Canada and involving more than six agencies (English et al. 2011).

Canada's main legal tool for sockeye salmon habitat conservation is the Fisheries Act, in place since 1976. It acknowledges the need to protect critical habitat for all life stages of the sockeye, during their food sources and the quality of water in which they live (Johannes et al. 2011). Again, it is overall a good measure that has been achieved through limitation of development, restoration of lost or damaged habitats and salmonid 'enhancement' in the form of hatcheries and spawning channels. Many habitat restoration and compensation projects have been ineffective, however (Wilson 2003), hatcheries have unintended, negative ecological impacts on the sockeye (see below, Kostow et al. 2009).

Escapement targets, set annually under the Pacific Salmon Treaty by an international panel, are complicated by the cyclic nature of Fraser River stocks and resulting interannual variability in return rates (English et al. 2011). Furthermore, measurement of actual escapement is complicated by the loss of sockeye (Figure 5). Escapement targets were not met for the Early Stuart sockeye from 2005–2009. While early targets met, English et al. (2011) conclude that overharvest likely occurred in the Early Stuart sockeye from 1984–2009 and for Early Summer sockeye from 1960–1989 (Figure 5).




Figure 5. Estimates of total, escapement and Eurotac losses for Fraser sockeye by return group. Eurotac losses were not estimated prior to 1992. English et al. 2011.

## Bristol Bay Management

English et al. (2011) reviewed differences in management structure between Bristol Bay and the Fraser River and made the following observations. The Fraser River is subject to a complex international management structure, management of the Bristol Bay sockeye falls

entirely within the Alaska Department of Fish Game and the Area Management Biologists. The simpler structure of Bristol Bay management allows for changes in the harvest regulations on day-to-day basis during the fishing season, while management decisions require a more lengthy process for Fraser River sockeye. Stock fishing issues are also related to the nature of the harvest fishing along the districts for inshore stocks, compared to the Fraser River with four runs timing groups which consist of more than 25 different stocks. There are limited inshore Bristol Bay stocks to the Fraser River. Furthermore, relatively escapement and low abundance populations in the Bristol Bay region, recreational and subsistence fisheries in Bristol Bay amount to less than 1% of the total harvest, while First Nations and recreational allocations are much higher in the more densely populated Fraser River. Bristol Bay's sockeye runs are also more concentrated, with a typical lasting six weeks compared to the more three months on the Fraser River. Over Bristol Bay benefits from a 'diversified portfolio' of stocks and life history types exploiting multiple large, productive rivers, resulting in extremely limited fisheries closures (Schindler et al. 2010). In contrast, Fraser River fisheries have been very limited or closed off the last 20 years.

Wide fluctuations in Bristol Bay's sockeye runs on the Fraser River (Figure 6) require managers to adjust goals every year, resulting in overharvest stocks. In the same period, variability in returns to Bristol Bay managers to cause fixed escapement based on the maximum sustained yield principles. Finally, escapement estimates in Bristol Bay are significantly more accurate than those in the Fraser River owing to the methodology where counts at sonar upstream of each commercial fishery in Bristol Bay essentially one hydroacoustic site in the Fraser River) and the fact that Bristol Bay fisheries are not subject to very high return mortality, to within (up to 90%) Fraser River stocks are subject to between migration and spawning grounds.

## Influences on Bristol Bay and Fraser River sockeye

Due to the economic importance and historically high returns, Bristol Bay and Fraser River sockeye salmon have been compared in recent months. Proponents of development in Bristol Bay and the Fraser River as example of major fisheries 'coexisting' (Joling 2011). However, despite watershed area of the Fraser Basin more than doubling Bristol Bay's — Fraser River sockeye abundance pales in comparison (Figure 6). Further, though the Kvichak River listing as a stock of concern (Mørstøl et al. 2010), Bristol Bay sockeye are not currently experiencing the types of decline exhibited in the Fraser River. Possible reasons for these differences abound, and are discussed below.

**Figure 6. Bristol Bay (blue) and Fraser River (red) total runs (catch + escapement) 1986.** Averages for each river are indicated by shading. Data from ADFG 2010 and PS6.

## Habitat

Bristol Bay encompasses nine major watersheds and has a drainage of 100,000 km<sup>2</sup> (FLBS 2011), while the Fraser River watershed drains 238,000 km<sup>2</sup> (Reynoldson et al. 2006). Further, watersheds are subject to opposite trends in productivity when Bristol Bay is experiencing low productivity, Fraser River and other U.S. west coast rivers experience lower productivity and vice versa (Mantua et al. 1997, Mantua and Hare 2002). The Bristol Bay basin was recently ranked as containing the most physically complex habitat throughout the range of Pacific salmon, making it a concern for future impacts of climate change (Mantua and Francis 2004, FLBS 2011) than most other Pacific watersheds supporting salmon, including the Fraser River (FLBS 2011).

## Aquaculture

In addition to management practices (reviewed by English 2011), it is important to note that major aquaculture activities in the Fraser River basin are in the straits, those in Bristol Bay where aquaculture is prohibited. About 10 fish are located on the migration route of Fraser sockeye salmon (Price et al. 2011). Although research is controversial, salmon have been associated with increased transmission of sea lice and disease (Price et al. 2011, Mitterer 2011). Globally, juvenile survival or abundance is reduced by aquaculture (Fordan and Myers 2008). A review

potential impacts to the Fraser sockeye from aquaculture activities is still forthcoming ([www.cohencommission.com](http://www.cohencommission.com)). □□□

Further, in response to population declines of the Fraser sockeye, the Canadian government operates nearly 30 hatcheries (MacDonald et al. 2011, Appendix 1). □ Unintended effects of the hatcheries include increased occurrence of disease (ischemia al. 2007), direct predation of fish by hatchery fish (Naiman and Sharpe 2001), □ competition for food resources (Dietman et al. 2011 in the past in freshwater environment (Tatara et al. 2008), in the estuary (Daly et al. 2011), and there is a (Ruggenberg et al. 2001). □ Effects of competition decrease productivity of salmon (Buhle 2009). □ British Columbia does not support any salmon hatcheries, and salmon farming is prohibited in the Bay and throughout the State of Alaska. □

## ***Human development*** □

More than two-thirds of British Columbians live in the Fraser River Basin with an overall population of 2.7 million residents in 2006 (BC 2010). Effects of human activities including urbanization, forestry, mining, agriculture, contaminant s, □ introduction s non-native species, and other factors are widely considered to be major factors in the declines of salmon worldwide (Nehlsen et al. 1991, □ Hartman et al. 2006 □ Bristol Bay currently supports only about seventeen small communities, and a population less than 5000 (DFO 2010). At present, the region does not support major industrial or other human activity. □ In contrast to water quality problems in the Fraser River discussed above, available data for waters in the Bristol Bay indicate cold, well oxygenated conditions with low concentrations of dissolved metals and other solutes (□ Zamzow 2000). □

## ***Cumulative impacts*** □

The analysis conducted for the Cohen Commission is limited to potential of the Fraser River sockeye declines within the past twenty years, during which declines became noticeable and commercially problematic (Pacific Salmon Commission data). □ Majority of experts released to date conclude that baseline and other pre-existing data is insufficient to do a thorough examination of the factors in question (Cook et al. 2004, Christensen and Tripp 2001, Hinch and Martin 2001, and others). □ Inquiry currently isolates individual potential factors in the declining fish population and synergistic effects of all factors combined. □ Christensen and Tripp (2001) conclude after the analysis of predation of sockeye on that "Cumulative threats are far more difficult to evaluate than a single factor. □ In case of Fraser sockeye salmon, stress from higher water temperatures, more in the competition due to the increased pressure with resulting lower growth, and the running the gauntlet through predators whose alternative prey may have diminished, may all have had cumulative effects. □ Assessing cumulative effects of these and other stresses will require integrated evaluation." □ □ Peter (2011) indicate that readers "should not necessarily expect to find single dominant cause of decline in the Fraser sockeye." □ Finally Healey (in press), in paper predicting Fraser sockeye response to climate change, indicates that the cumulative impacts of climate across life stages

will be much greater than the impacts on individual species. He concluded that the impacts will also carry forward to the ecosystem, potentially leading to a downward spiral of the productive capacity," predicting a dark future for Fraser River sockeye salmon not unlike that of other major salmon rivers south of along the Pacific coast where salmon are extirpated from 40% of their former range (NRC 1996).

## Conclusions

Fraser River sockeye salmon populations are suffering from a host of myriad problems associated with urban and industrial development, leading to dramatic decreases in productivity, multiple fisheries closures, and federal and international population listings. In the freshwater environment from mining, wood product and other industrial facilities, wastewater treatment plants, landfills, and salmon enhancement facilities (e.g., hatcheries and spawning channels) has led to contamination of over 500 sites, causing problems with pH, TSS, turbidity, nutrients, metals, phenols, personal care products, and pharmaceuticals. Introduced predators such as the yellowtail smelt, as well as the well-known hatchery smolt impact Fraser River sockeye in the freshwater environment. And finally, increased river temperatures resulting from climate change are associated with higher mortality of the sockeye smolt at spawning grounds likely due to increased physiological stress at higher temperatures, decreased swimming efficiency, and faster development of the pathogens.

In the marine environment, industrialization and urban growth has led to the contamination of the Strait of Georgia by polybrominated diphenyl ethers, personal care products, and pharmaceuticals. Dredging likely has reduced marine and estuarine sockeye habitat. Increased ship traffic is associated with accidental spills, noise, and introduction of non-native species. Warmer temperatures resulting from climate change are associated with more frequent harmful algal blooms, resulting in low oxygen levels in the marine environment, as well as decreased plankton levels which are an important source of salmon food source.

Current efforts to understand sockeye declines isolate potential causes, failing to consider the synergistic effects of the combination of such as the contamination and use of introduced predators, climate change, and others. Further analyses are needed to rely upon inadequate historical datasets, which fail to satisfactorily baseline conditions.

Given the distinct physical and biological nature, as well as higher levels of urbanization and industrialization in the Fraser River basin relative to Bristol Bay basin, recent comparisons between the two watersheds are suspect. However, when comparing sockeye salmon populations alone, Bristol Bay — the world's largest sockeye salmon producing system — outnumbers the Fraser River by 10 times in watershed less than half its size. Indeed, the comparison between the two systems may simply highlight the inability of the human component to coexist with salmon.

## References □ð

- ð  
Angelo, M., P. Gallagher, J. Orr, R. Reynolds, D. Richardson, K. Wilson. 2010. Fraser sockeye □ð 2010: Findings of student □ðsts' think tank. □ð Speaking Salmon. 6 pp. □ð
- ð  
Batten, S. Dean, D. L. MacKas. 2009. Shortened duration of □ð annual □ð *Neocalanus plumchrus* □ð biomass peak in □ð Northeast Pacific. Marine Ecology Progress Series 393: 189 – 198. □ð
- ð  
Beamish, R.J., D.J. Noakes, G.A. McFarlane, L. □ð Klyashtorin, V. Ivanov, and V. □ð Kurashov. 1999. The regime concept and natural trends in □ð productivity of □ð Pacific salmon. □ð Canadian Journal of □ð Fisheries and Aquatic Sciences 56(8): 516 – 526. □ð
- ð  
Beamish, R.J. and C.M. Neville. 2001. Predation □ð base □ð mortality on □ð juvenile salmonids in □ð Strait of □ð Georgia. North Pacific Anadromous Fish Commission (NPAFC) Bulletin 2: 11 – 13. □ð
- ð  
Beauchamp, D.A. 1995. Riverine production on □ð sockeye salmon migrating to □ð Lake □ð Washington. North American Journal of □ð Fisheries Management 15(2): 353 – 365. □ð
- ð  
Bradford, M.J., C.P. Torrey, and L.M. □ð Herbig. 2008. Biological risk assessment for yellow perch (*Percula flavescens*) □ð British Columbia. Canadian Science Advisory Secretariat. Research Document 2008/076, 23 pp. □ð
- ð  
Bradford, M.J., J. □ð Vliet, and D.A. Patterson. 2000. Infection of □ð anguilicidone of □ð Fraser □ð sockeye salmon, *Oncorhynchus nerka* (Walbaum), by *Parvicapsula minibicornis* and its □ð effect on □ð physiology. Journal of □ð Disease 33: 69 – 77. □ð
- ð  
Buhle, E.R., K.K. Hobson, M.B. Scheuerell, and A. □ð Alba. 2009. Using □ð unplanned □ð experiment to □ð evaluate effects of □ð hatchery environmental variation on □ð threatened □ð populations of □ð salmon. Biological Conservation 140: 2449 – 2455. □ð
- ð  
Burgner, R.J. 1991. Life history of □ð sockeye salmon (*Oncorhynchus nerka*). □ð pp. 112. □ð  
Groot, C. □ð Land, D. Margolis. (Eds.). Pacific Salmon Life Histories. □ð PBS, Vancouver, British Columbia. □ð
- ð  
Cartwright, M.A., D.A. Beauchamp, and M.D. Bryant. 1998. Quantifying cutthroat trout □ð (*Oncorhynchus clarkii*) predation on □ð sockeye salmon (*Oncorhynchus nerka*) □ð using a □ð bioenergetics approach. Canadian Journal of □ð Fisheries and Aquatic Sciences 55: 1285 – 1295. □ð
- ð  
Chandler, P. □ð 2010. □ð temperature and salinity at □ð big holes. Pp. 49–51. □ð  
Crawford, and J. Irvine (Eds.). State of □ð physical, and selected fishery resources □ð of □ð Canadian marine ecosystems in □ð 2009. □ð
- ð  
Chittenden, C.M., R.J. Beamish, and R.S. McKinley. 2009. A □ð critical review of □ð Pacific □ð marine research relating to □ð climate □ð Journal of □ð Marine Science 66: 2195 – 2224. □ð
- ð

- Christensen, V. and Trides. 2001. Predation on the Fraser River sockeye salmon. The Cohen Commission of the Inquiry into the Decline of the Sockeye Salmon in the Fraser River. 130 pp. □
- 
- Clearly, J.S., J.E. Schweiger, and V. D. H. 2009. Stock assessment and management advice for the British Columbia herring fishery: 2009 assessment and 2010 forecasts. Canadian Science Advisory Secretariat Research Document 2009/079. 89 pp. □
- 
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2003. COSEWIC assessment and status report on the sockeye salmon (*Oncorhynchus nerka*) (Coho population) in Canada. 8 pp. □
- 
- Cooke, S.J., S.G. Hinrich, A.P. Farrell, M.J. Lapointe, S.R. M. Jones, J.S. Macdonald, D.A. Patterson, M.O. Haley, and G. D. Van Kraan. 2004. Abnormal migration timing and high en route mortality of the sockeye salmon in the Fraser River, British Columbia. *Fisheries* 29(2): 22-33. □
- 
- Cosgrove, J.A. 2005. The first specimens of the Humboldt in the British Columbia. PICES Press. July: 30-31. □
- 
- Crawford, W.H. and J.R. Urquhart. 2010. State of the physiological and selected fishery resources of the Pacific marine ecosystems in the Department of the Fisheries and Oceans Canadian Science Advisory Secretariat. Research Document 2010/053. 137 pp. □
- 
- Crossin, G.T., S.G. Hinrich, S.J. Cooke, M.S. Cooperman, D.A. Patterson, D.W. Welch, K.G. Hanson, I. Olszynski, K. English, and A.P. Farrell. 2009. Mechanisms influencing the timing and success of the reproductive migration in Capital of the breeding semelparous fish species, the sockeye salmon. *Physiological and Biochemical Zoology* 82(6): 635-652. □
- 
- Daly, E.A., R.D. Brodeur, J.P. Fisher, L.A. Westkamp, D.J. Reed, and B.R. Beckman. 2001. Spatial and trophic overlap of marked and unmarked Columbia River spring Chinook salmon during early marine residence with implications for competition between hatchery and naturally produced fish. *Environmental Biology of Fishes* published online: 04 June 2011. □
- 
- DCRA (Division of the Community Regional Affairs). 2010. Community Information Summaries (CIS). Alaska Community Database. [http://www.dced.state.ak.us/dca/commdb/cf\\_cis.htm](http://www.dced.state.ak.us/dca/commdb/cf_cis.htm). □ Access date: 11 Aug 2011. □
- 
- DFO (Fisheries and Oceans Canada). 2003. Annual sockeye salmon stock status report. Fisheries and Oceans Canada, Pacific Region. 60 pp. □
- 
- DFO. 2001. Commercial catch statistics. [http://www.pac.dfo-mpri.gc.ca/stats/comm/index\\_english.htm](http://www.pac.dfo-mpri.gc.ca/stats/comm/index_english.htm). Accessed 10 May 2011. □
-

- Dittman, A.B. & D. May, M. Larsen, M. Johnstone, D. Fast, In the press. Spatial and temporal overlap of hatchery and wild spring Chinook salmon spawning: Effects of hatchery accumulation sites. *Environmental Biology of Fishes*. □
- Eggers, D.M. and J.R. Dryne. 2007. Trends in abundance and biological characteristics for North Pacific sockeye salmon. North Pacific Anadromous Fish Commission (NPAFC) Bulletin 4: 1-73. □
- El-Sabaawi, R., P.E. Dover, M. Kainz, A. Mazum. 2009a. Characterizing dietary variability and trophic position of coastal copepods: insights into leiosomes and fatty acids. *Marine Biology* 156(3): 225 - 237. □
- El-Sabaawi, R., P.E. Dover, M. Kainz, A. Mazum. 2009b. Interannual variability in the fatty acid composition of copepods *Neocalanus plumchrus* in the Strait of Georgia, British Columbia. *Marine Ecology Progress Series* 382: 153 - 164. □
- Eliason, E.J., T.D. Clack, M.J. Hage, L.M. Hanson, Z.S. Gallagher, K.M. Jeffries, M.K. Gale, D.A. Patterson, S.G. Hinrichs, and A.P. Farrell. 2001. Differences in thermal tolerance among sockeye salmon populations. *Science* 330: 109 - 112. □
- Ellis, E.M. & D. Church. 2004. Characterisation of secondary channels of the lower Fraser River. Habitat Conservation Trust Final Report. 188pp. □
- English, K.K., D. Egell, R.C. Bocking, M.K. Link, and S.W. Raborn. 2001. Fraser River sockeye fisheries and fisheries management and comparison with Bristol Bay sockeye fisheries. The Columbia Commission of the Inquiry into the Decline of Sockeye Salmon in the Fraser River. □ 317pp. □
- Environment Canada. 2007. British Columbia and Yukon Territory water quality report (2001-2004). Application of Canadian Water Quality Index. Environment Canada. 828 pp. □
- Errington, J.C. and K.D. Ferguson. □ 1987. Acid rain damage in British Columbia: Today and tomorrow. Proceedings of the Annual British Columbia Mine Reclamation Symposium in Kamloops, BC 1987. Technical and Research Committee on Mine Reclamation. □ pp23 □
- Farrell, A.P., S.G. Hinrichs, S.J. Cooke, D.A. Patterson, G.T. Crossin, M. Lapointe, M. Mathes. 2008. Pacific salmon in water: Applying biotelemetry models and biotelemetry to predict success of spawning migrations. Physiological and Biochemical Zoology 81(6): 693 - 707. □
- Fayram, A.H. and T.H.S. Hobley. 2000. Impact of predation by smallmouth on sockeye salmon in Washington, Washington. North American Journal of Fisheries Management 20: 81-89. □

- FBC(Fraser Basin Council). 2000.The Fraser: A 300 km long river. Fraser Basin Council in 300 collaboration with the BC Ministry of Environment 300 pp. 300
- FLBS (Flathead Lake Biological Station). 2001.Riverscape Analysis Project. FLBS, 300 University of 300 Montana <http://rap.ntsg.umt.edu/home>. 300 300 Accessed August 2001. 300
- Foote, C.J., and G.S. Brown. 1998.Ecological relationship between freshwater sculpins 300 (genus *Cottus*) 300 and spawning sockeye salmon (*Oncorhynchus nerka*) 300 in Alaska., 300 Canadian Journal of 300 Fisheries and Aquatic Sciences 55(15):1524 – 1536. 300
- Ford, J.S. and R.A. Myers. 2008. A global assessment of salmon aquaculture impacts on 300 wild salmonids. 300 Biology 8(2):e33 DOI:10.1371/journal.pbio.0060033. 300
- Forrest, K.W., J.D. Cave, C.G. Michielsens, M. Hauke, and D.M. Smith. 2009. Evaluation of 300 an 300 effective 300 detection 300 of 300 salmonid 300 gill net 300 fisheries. North 300 American Journal of 300 Fisheries Management 29(4):885 – 894. 300
- FREMP (Fraser River Estuary Management Plan). 2006. Environmental management 300 strategy for dredging in the Fraser River Estuary. 470 300 pp. 300
- Gallaugh, P. 300 Land 300 Water 2010. Summit on 300 Fraser sockeye salmon: 300 Understanding stock declines and prospects for the future. Speaking for the Salmon. 210 300 pp. 300
- Grant, S.C.H., C.G. Michielsens, E.J.P. Ross, and A. 300 2010. Predicting run size forecasts 300 for Fraser River sockeye salmon in 300 the 2010 Department of 300 Fisheries and Oceans Canadian 300 Science Advisory 300 Secretariat. Research Document 2010/042, 120 300 pp. 300
- Groot, C. 300 Land 300 March 1991. Pacific salmon life 300 fe 300 historical 300. 560 300 pp. 300
- Hague, M.J., M.R. Feit, J.R. Miller, D.A. Patterson, G.L. Russell, A.P. Fadell, and S.G. Hinch. 300 2011. Modeling the future hydroclimatology of 300 lower Fraser River and its impacts on 300 the spawning migration survival of 300 sockeye salmon. Global Change Biology 17:870 – 980. 300
- Hare, S., R.N. Mantua, and R.C. Francis. 1999. Inverse production regimes: 300 Alaska 300 West Coast Pacific salmon. Fisheries 24(1):60 – 14. 300
- Hartman, G.F., T. Connorthcote, and C.J. Cederholm. 2006. Human numbers: the alpha factor 300 affecting the future of 300 salmon. 300 pp. 260 – 290. 300 Salmon 2100: 300 Future of 300 Pacific 300 Salmon, R.T. Lackey, D.H. Ladd, and S.L. Duncan (Eds.). 300 American Fisheries Society, 300 Bethesda, MD. 300
- Hinch, S.G. and E.G. Martins. 2001. A 300 review of 300 potential climate change effects on 300 survival 300 of 300 Fraser sockeye salmon and an 300 analysis 300 interannual trends in 300 the loss and 300 pre-spawning mortality. The Columbia Commission of 300 the Inquiry into the Decline of 300 Sockeye salmon 300 in 300 the Fraser River. 110 300 pp. 300

- ③
- IPCC (International Panel on Climate Change). 2007. Climate Change 2007: Synthesis Report. Contribution of Working Group II to the Fourth Assessment. 104 pp. □ ③
- Johannes, M.J.S., L.H.Niko, R.J.R. Hoogendoorn, and R.E.Scott. 2011. Fraser River watershed key habitat uses in the Lower Fraser and Strait of Georgia. Geofish Commission of the Inquiry into the Decline of Sockeye Salmon in the Fraser River. 116 pp. □ ③
- Johannessen, S.C. and R.W. McDonald. 2009. Effects of long-term climate change on fish and sea: Threats of the Geographical Columbia, Canada. Climate Research 40(1): 1-21. □ ③
- Joling, D. 2011. Executive says Pebble, fishing camp could exist. Grist. 04 Mar 2011. □ ③
- Kistritz, R.L. 1994. Habitat compensation, restoration, and creation in the Fraser River estuary: Achieving net ecological habitat. Canadian Manuscript Report of the Fisheries and Aquatic Sciences. 118 pp. □ ③
- Kostow, K. 2009. Factors that contribute to ecological risks of salmonid hatchery programs and some mitigating strategies. Reviews in Fish Biology and Fisheries 19(1): 9-31. □ ③
- Levin, P.S. & R.W. Zabel, and J.G. Williams. 2001. The road to extinction paved good intentions: negative interaction of fisheries with threatened salmon. Proceedings of the Royal Society of London B Biological Sciences 268: 1153-1158. □ ③
- Levings, C.D. and D.J.M. Nishimura. 1997. Created and restored marshes in the Lower Fraser River, British Columbia: summary of their functioning as fish habitat. Water Quality Research Journal of Canada 32: 9-18. □ ③
- Luck, M.A. & D. Maumellee, D. Whited, D. Lusk, D. Chilcott, D. Borlang, D. Goodwin, D. McDonald, J. Kimball, J. Stan. 2010. Remote sensing analysis of physical complexity of Pacific Rim rivers to assist salmon conservation. Earth Surface Processes and Landforms 35: 1330-1343. □ ③
- MacDonald, D.J. & D. Sinclair, D. Crawford, D. Prentice, M. Meneghini. 2011. Potential effects of contaminants on river sockeye salmon. The Cohen Commission of the Inquiry into the Decline of Sockeye Salmon in the Fraser River. Technical Report 2016-1 and appendices. □ ③
- Mantua, N.J. & S.R. Hare, Y.Q. Zhang, M. Wallace, and R.C. Francis. 1997. A Pacific decadal climate oscillation with impacts on salmon production. Bulletin of the American Meteorological Society 78(6): 1069-1079. □ ③
- Mantua, N.J. and S.R. Hare. 2002. The Pacific Decadal Oscillation. Journal of Oceanography 35(1): 35-44. □ ③

- ð  
Mantua, N.J. and R.C. Francis. □ð 2004. National insurance for Pacific northwest salmon and salmon fisheries: □ð finding a way through the tangled bank. □ð In E.E. Knudsen □ð and D. MacDuff (Eds.), □ð Sustaining Management of □ð North American Fisheries. □ð American □ð Fisheries Society Symposium 43:123 - 147. □ð
- ð  
Martell, S.J.D., C.J. Waters, □ð and R. Hilborn. □ð 2008. Retrospective analysis of □ð harvest management performance for Bristol Bay and Fraser River sockeye salmon (□ð *Oncorhynchus nerka*). □ð Canadian Journal of □ð Fisheries and Aquatic Sciences 65:409 - 424. □ð
- ð  
Miller, K.M., S. Okiishi, J. Kaukinen, N. Gither, E. Ham, M.R. Curtis, D.A. Patterson, T. □ð Sierociński, L. □ð Donnison, □ð P. S. Hinch, K. A. Hruska, S.J. Cooke, K. □ð English, and A.P. □ð Farrell. □ð 2011. Geosignatures predict migration and spawning failure in □ð wild □ð Canadian salmon. □ð *Science* 331:214 - 216. □ð
- ð  
MOE (British Columbia Ministry of □ð Environment). □ð 1985. Ambient water quality assessment □ð and objectives for the Fraser River upstream from Kamloops to the Ministry of □ð Environment, Government of □ð British Columbia. □ð
- ð  
MOE. □ð 2006. B.C. □ð municipal waste tracking report. British Columbia Ministry of □ð the Environment. 100pp. □ð
- ð  
MOE. □ð 2008. Environmental trends in □ð British Columbia: □ð 2008 British Columbia Ministry of □ð the Environment. 360pp. □ð
- ð  
Morstad, S.M. □ð Jones, □ð Sands, □ð Salomone, □ð Baker, □ð Bank, □ð West. □ð 2009. □ð Bristol Bay area annual management report. ADFG, Division of □ð Fisheries and Division of □ð Commercial Fisheries. 140pp. □ð
- ð  
MPCA (Minnesota Pollution Control Agency). □ð 2008. Endocrine disrupting compounds: □ð A □ð report to the Minnesota Legislature. 340 □ð pp.
- ð  
Muir, D.C.G., R.C. Shearer, J. □ð Vandam, S.C. Donaldson, and C. □ð Fu. □ð 2015. □ð Contaminants in □ð Canadian biota and implication for human health: □ð Conclusions □ð knowledge gaps. □ð *Science of the Total Environment* 535:39 - 54. □ð
- ð  
Naish, K.A., J.E. Taylor III, □ð S. Levan, T.P. Quinn, J.R. Winton, D. □ð Huppert, □ð R. □ð Hilborn. □ð 2007. An □ð evaluation of □ð the effects of □ð conservation and fishery enhancement hatcheries on □ð wild populations of □ð salmonids in □ð the Marine □ð gy 1653-197. □ð
- ð  
Naman, S. □ð Casals, □ð. □ð 2011. Predation by □ð hatchery yearling salmonids on □ð wild □ð subyearling salmonids in □ð the freshwater environment: □ð A □ð review with □ð case histories, and implications for □ð management. □ð *Environmental Biology of □ð Fishes*. □ð Published □ð online: 27 □ð □ð April. □ð □ð

- Nehlsen, W., J.E. Williams, and J.A. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk California, Oregon, Idaho, and Washington. *Fisheries* 16(2):40–72. [\[Full Text\]](#)
- Nelitz, M., M. Poeter, Parkin Kon, W. Wieckowski, D. Marmotek, B. Bryan, and D. Abraham. 2001. Evaluating the status of Fraser River sockeye salmon and role of the freshwater ecology in the decline. The Canadian Commission of Inquiry on the Decline of Sockeye Salmon in the Fraser River. 2001:36 pp.
- NRC (National Research Council). 1996. Upstream: salmon and society in the Pacific Northwest. NRC Committee on Protection and Management of the Northern Pacific Northwest Anadromous Salmonids, Board on Environment Studies and Toxicology, Commission on Life Sciences. 450 pp.
- Olesiuk, P.F. 2009. Abundance of Steller sea lions (*Eumetopias jubatus*) in British Columbia. Canadian Science Advisory Secretariat: Research document 2008/063, 33 pp.
- Pestal, G. and A. Andrade. 2009. Using qualitative risk evaluations to prioritize resource assessment activities for Fraser River sockeye. Canadian Scientific Advisory Secretariat: Research Document 2009/071, Fisheries and Oceans Canada. 81 pp.
- Peterman, R.M. and B. Domon. 2011. Fraser River sockeye production dynamics. The Canadian Commission of Inquiry on the Decline of Sockeye Salmon in the Fraser River. Technical Report 10, 133 pp.
- Peterman, R.M. and F.Y.C. Wong. 1994. Cross correlations between reconstructed ocean abundances of British Columbia sockeye salmon (*Oncorhynchus nerka*). Canadian Journal of Fisheries and Aquatic Sciences 51:1814–1824. [\[Full Text\]](#)
- Peterman, R.M., D. Marmotek, B. Beckman, B. Bradford, M. Baird, and M. Bell. 2001. Scheuerell, M. Staley, W. Wieckowski, K. Winson, and C. Wood. 2000. Synthesis of evidence from a workshop on the decline of Fraser sockeye. A report of the Pacific Salmon Commission, Vancouver BC. 240 pp.
- Petersen, J.H. 2001. Density, aggregation, and body size of the northern anchovy preying on juvenile salmonids in the Fraser River. Journal of Fish Biology 58:1137–1148. [\[Full Text\]](#)
- Price, M.H.H., S.L. Luboszcz, R.D. Routledge, A.S. Gottesfeld, C. Bjørn, M. Reynolds, and R. Reynolds. 2001. Sea louse infection of juvenile salmonids in relation to marine farm salmon in Canada's west coast. PLoS One 6(2):1–9. [\[Full Text\]](#)
- Rand, P.S. 2008. *Oncorhynchus nerka* (Fraser River). In: IUCN Red List of Threatened Species, [www.iucnredlist.org](http://www.iucnredlist.org). Version 2010.4. [\[Full Text\]](#)
- Rand, P.S., C.P. Kelton, X. Augerot, G. Rijksen, and G.T. Ruggerone. 2007. Comparison of sockeye salmon (*Oncorhynchus nerka*) monitoring Fraser River Basin, 2000–2005.

British Columbia, Canada and Bristol Bay, Alaska, USA. North Pacific Anadromous Fish Commission (NPAFC) Bulletin 4: 27–28. 2018. 27 pp. G

- Rand, P., S. G. Hinch, J. M. Morris, G. Foreman, M. J. McDonald, J. S. McDonald, M. O. Healey, A. P. Farrell, D. A. Higgs. 2006. Effects of discharge, temperature, and future climates on the energetics and mortality of adult migrating Fraser River sockeye salmon. *Transactions of the American Fisheries Society* 135: 655–667. 2018. 13 pp. G
- Rensel, J.E., N. D. Haight, T.J. Flynn. 2010. Fraser River sockeye salmon marine survival decline and harmful blooms of *Heterosigma akashiwo*. *Canadian Journal of Zoology* 88(1): 98–111. 2018. 14 pp. G
- Reynoldson, T.B., J. D. Culp, D. Lovewell, and R. Richardson. 2005. Fraser River Basin. Pp. 96–732 in *Rivers of the Northern Hemisphere*. A. C. Benke, A. Canfield, and C. E. Cushing (Eds.). Elsevier Academic Press, Burlington, MA. 2018. 13 pp. G
- Roby, D.D., D.E. Lyons, D.P. Crabb, K. D. Cahill, G.H. Visser. 2003. Quantifying the effect of predators on the endangere species of salmonids bioenergetics approach: the Castries and the juvenile salmonids in the Columbia River estuary. *Canadian Journal of Zoology* 81(2): 26–36. 2018. 11 pp. G
- Ruggerone, G. T. and D. L. Nielsen. 1992. Predation on the sockeye salmon fry by juvenile chinook salmon in the Chignik Lakes, Alaska: implications for salmon management. *North American Journal of Fish Management* 12(3): 10–11. 2018. 2 pp. G
- Ruggerone, G. T. and D. L. Nielsen. 2004. Evidence for competitive dominance of the pink salmon (*Oncorhynchus gorbuscha*) over other salmonids in the North Pacific Ocean. *Reviews in Fish Biology and Fisheries* 14(3): 37–39. 2018. 3 pp. G
- Ruggerone, G. T., B. A. Agler, and D. L. Nielsen. 2011. Evidence for competition at the seaweed Norton Sound chum salmon and Asian hatchery chinook salmon environmental Biology of Fishes. Published online: 07.02.2011. 2018. 12 pp. G
- Schindler, D.E., P.R. Leavitt, C.S. Brock, S.P. Johnson, and P.D. Quinn. 2005. Marine derived nutrients, commercial fisheries, and production of the salmonid algae in the Alaska. Ecology 86(12): 3205–3221. 2018. 16 pp. G
- Schindler, D.E., R. D. Hill, B. Chas-Poole, R. Wright, T.P. Quinn, L.A. Rogers, and M.S. Webster. 2010. Population diversity and the portfolio effect in exploited species. *Nature* 465: 609–612. 2018. 4 pp. G
- Stewart, I.T., D.R. Cayelan, and M. D. Dettling. 2005. Changes toward earlier streamflow timing across Western North America. *Journal of Climate* 18(11): 2714–2723. 2018. 10 pp. G
- Tatara, C.P., S.C. Riley, and J.A. Schaeurer. 2008. Environmental enrichment in the steelhead (*Oncorhynchus mykiss*) hatcheries field valuation of the aggression and territoriality in the natural hatchery fry. *Canadian Journal of Fish and Aquatic Sciences* 65(7): 744–753. 2018. 10 pp. G

□ð

Tovey, C.P., M.J. Bradford, and L.D.M. G. Herb. 2008. Biological risk assessment for smallmouth bass (*Micropterus dolomieu*) and largemouth bass (*Micropterus salmoides*) in British Columbia. Canadian Science Advisory Secretariat Research Document 2008/075. 47 pp.

□ð

Veldhoen, N., M.G. Ikonomou, C. Duletz, D. MacPherSon, S. Sampson Kelly, and C.C. Helbing. 2010. Gene expression profiling and environmental contaminant assessment of migrating Pacific salmon in the Fraser River watershed of British Columbia. Aquatic Toxicology 97:212 - 225. □ð

□ð

Wagner, G.N., S.G. Hinrich, L.J. Kuchel, A. L. L. R. M. Jones, D.A. Patterson, J. McDonald, G. Van Der Kraak, M. Shrimpton, K. English, S. Laishram, M. Healey, and A.P. Farrell. 2005. Metabolic rates and swimming performance of adult Chinook salmon (*Oncorhynchus nerka*) after control feeding with *Pavicapsula miliabicornis*. Canadian Journal of Fisheries and Aquatic Sciences 62:2124 - 2133. □ð

□ð

Warren, Jr., M.L. and M.C. Pandew. 1998. Roads as barriers to small stream fish movement. Transactions American Fisheries Society 127:630 - 644. □ð

□ð

Wilson, P.H. 2003. Using population projection matrices to evaluate recovery strategies for the Snake River spring and summer Chinook salmon. Conservation Biology 17(8):782 - 794. □ð

□ð

Winship, A.J. and A.W. Trippes. 2003. Prey consumption of Stellar sea lions (*Eumetopias jubatus*) in Alaska: how prey do they intend? Fishery Bulletin 101(1):143 - 161. □ð

□ð

Zamzow, K. 2011. Surface water quality near the proposed Pebble Mine, Alaska, 2009 - 2010: Nushagak, Kvichak, and Chilikna drainage headwaters. Center for Science in the Public Participation. □ð

**Appendix 1. Salmonid Enhancement facilities in the Fraser River Basin. Not Available. McDonald et al. 2011**

Area of Interest/Facility Name	Facility Type	Species Targeted	Organization
<b>Cultus Lake</b>			
Chilliwack River Hatchery	Hatchery	Chinook, Coho, Chum, and Steelhead	DFO Operations
Fraser Valley Trout Hatchery Centre Creek Streamkeeper Program	Hatchery Hatchery	Native and Domestic Rainbow Trout, Anadromous and Coastal Cutthroat Trout, and Steelhead Trout N/A	Freshwater Fisheries Society of BC Public Involvement Programs (Volunteer)
<b>Harrison River</b>			
Chehalis River Hatchery Weaver Creek Spawning Channel Fee Creek Spawning and Rearing Channel	Hatchery Spawning Channel Hatchery	Coho, Chinook, Chum, Steelhead and Cutthroat Trout Sockeye, Chum, Pink Coho	DFO Operations DFO Operations Public Involvement Programs (Volunteer)
<b>Lower Fraser River</b>			
Inch Creek Hatchery Bell-Irving Kanaka Creek Hatchery Beecher Creek Streamkeepers Al Grist Memorial Hatchery Chilliwack River Action Committee (Trap Site) Stave Valley Salmonid Enhancement Society Nicomen Slough Spawning Channel Musqueam Creek Project Steveston High School Hatchery (on-site) Cougar Creek Salmonid Enhancement Group Hoy Creek Hatchery River Springs Salmon Enhancement and Stream keepers	Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery Hatchery	Coho, Chinook, Chum, and Steelhead Trout Chum, Coho, Pink, Steelhead, and Cutthroat Trout Coho, Cutthroat, and Rainbow Trout Coho, Chinook, and Pink Steelhead Trout, Coho, Chinook, Chum, and Pink Coho and Chum Coho and Chum Coho, Chum, and Cutthroat Trout Coho and Chinook Coho Coho	DFO Operations Public Involvement Programs (Volunteer) Public Involvement Programs (Volunteer)

<b>Area of Interest/Facility Name</b>	<b>Facility Type</b>	<b>Species Targeted</b>	<b>Organization</b>
<b>Lower Thompson River</b>			
Spius Creek Hatchery	Hatchery	Chinook, Coho, and Steelhead Trout	DFO Operations
Loon Creek Hatchery	Hatchery	Rainbow Trout and Kokanee	Community Development Program Hatcheries
Deadman River Hatchery	Hatchery	Chinook and Coho	Community Development Program Hatcheries
<b>Nechako River</b>			
Nadina River Spawning Channel	Spawning Channel	Sockeye	DFO Operations
Spruce City Wildlife Fish Hatchery	Hatchery	Chinook	Public Involvement Programs (Volunteer)
<b>North Thompson River</b>			
Clearwater Trout Hatchery	Hatchery	Rainbow Trout and Kokanee Salmon	Freshwater Fisheries Society of BC
Dunn Lake Hatchery	Hatchery	Coho and Chinook	Community Development Program Hatcheries
<b>Pitt River</b>			
Upper Pitt River Hatchery	Hatchery	Chinook and Sockeye	DFO Operations
ALLCO Hatchery	Hatchery	Coho, Steelhead, Cutthroat, Pink, and Chinook	Public Involvement Programs (Volunteer)
Hyde Creek Hatchery	Hatchery	Coho and Chum	Public Involvement Programs (Volunteer)
<b>Quesnel River</b>			
Horsefly Spawning Channel	Spawning Channel	Sockeye	DFO Operations
<b>Seton -Portage</b>			
Gates Creek Spawning Channel	Spawning Channel	Pink	DFO Operations
Seton Creek Spawning Channels	Spawning Channel	Pink	DFO Operations
<b>South Thompson River</b>			
Shuswap River Hatchery	Hatchery	Chinook	DFO Operations
Kingfisher Community Hatchery	Hatchery	Coho, Spring, Sockeye, and Kokanee	Public Involvement Programs (Volunteer)
Adams River	Fishway	Sockeye	DFO Operations

□ Ø

Area of Interest/Facility Name	Facility Type	Species Targeted	Organization
<b>Upper Fraser River</b>			
Penny Hatchery	Hatchery	Chinook	Community Development Program Hatcheries
Anderson Lake Fish Hatchery	Hatchery	Sockeye and Kokanee	Public Involvement Programs (Volunteer)
Hells Gate Fishways	Fishway	Sockeye, Coho, Pink, Chinook, Steelhead Trout	DFO Operations

□ Ø